

Instructions for Use

Bordered
SNS77733-3"x3"
SNS77744-4"x4"

Non-Bordered
SNS74502-2"x2"
SNS74520-4"x5"

Silicone Foam Wound Dressing

Simpurity™ Silicone Foam Wound Dressings contain a highly absorbent foam pad and gentle adhesive silicone contact layer, available with or without an adhesive border. These dressings help maintain a moist wound healing environment with no irritation upon removal, minimizes trauma to the wound and reduces pain during dressing changes.

Application:

1. Clean the wound with sterile cleanser or saline water and dry the surrounding skin
2. Choose the right size of dressing to cover the wound according to wound area

Bordered Silicone Foam:

1. Remove liners from the back side of the dressing
2. Remove one of the back layers and apply that dressing half to the wound. Then remove the second liner and secure the rest of the dressing to the wound

Non-Bordered Silicone Foam:

1. Remove liners from the back of the dressing
2. Apply the dressing to the wound bed
3. Secure the dressing with medical tape or a transparent dressing

Changing Frequency:

Dressing change frequency will depend on the patient's condition and the level of exudate. This dressing is recommended to be changed three (3) times a week.

Dressing Removal:

Gently remove the tape, if used. Peel from the edge of the used foam dressing to remove it from the wound bed. Repeat application steps.

Indications:

Simpurity™ Silicone Foam dressings are indicated for the management of moderate to heavily exuding wounds such as: pressure ulcers, leg ulcers, non-infected diabetic foot ulcers, donor sites, post-operative wounds, lacerations and abrasions.

Contraindications:

The product is not indicated for third-degree burns or surgical implantations.